

Tsemo Aristide
 The International Center for Theoretical Physics
 Strada Costiera, 11
 Trieste, Italy.
 email tsemo@ictp.trieste.it

New Address:
 Tsemo Aristide,
 3738, Avenue de Laval Appt 106
 Montreal Canada H2X 3C9
 email: tsemoaristide@hotmail.com

Closed similarity lorentzian affine manifolds.

Abstract.

A $Sim(n-1,1)$ affine manifold is a n -dimensional affine manifold whose linear holonomy lies in the similarity Lorentzian group but not in the Lorentzian group. In this paper, we show that a compact $Sim(n-1,1)$ affine manifold is incomplete. Let $\langle, \rangle_L$ be the Lorentz form, and q the map on $\mathbb{R}^n$ defined by $q(x) = \langle x, x \rangle_L$. We show that for a compact radiant $Sim(n-1,1)$ affine manifold M whose developing map is injective, if a connected component C of $\mathbb{R}^n - q^{-1}(0)$ intersects the image of the universal cover of M by the developing map, then either C or a connected component of $C - H$, where H is an hyperplane is contained in this image.

Introduction.

An n -dimensional affine manifold M , is an n -dimensional differentiable manifold endowed with an atlas whose coordinate changes are locally affine maps. The affine structure of M pulls back to its universal cover $\hat{M}$, and defines on it an affine structure determined by a local diffeomorphism $D : \hat{M} \rightarrow \mathbb{R}^n$, called the developing map. The developing map gives rise to a representation $h : \pi_1(M) \rightarrow Aff(\mathbb{R}^n)$, called the holonomy of the affine manifold. Its linear part $L(h)$, is called the linear holonomy of the affine manifold. We will say that the affine manifold is complete, if and only if the developing map is a diffeomorphism. An n -affine manifold is said to be radiant if its holonomy fixes an element of $\mathbb{R}^n$.

We denote by $O(p, q)$, the subgroup of linear automorphisms of $\mathbb{R}^n$ which preserve a bilinear symmetric form of type p, q , and by $Sim(p, q)$ the group generated by $O(p, q)$ and the homotheties. An $O(p, q)$ affine manifold M is an affine manifold M such that the image of its linear holonomy $L(h)$ is a subgroup of $O(p, q)$. An $Sim(p, q)$ affine manifold M is an affine manifold M such that the image of its linear holonomy $L(h)$ is a subgroup of $Sim(p, q)$, and contains an element which is not in $O(p, q)$.

Let consider the flat riemannian torus T^n , Bieberbach has shown that closed $O(n, 0)$ affine manifolds are finitely covered by T^n . Using the notion of

discompacity, Yves Carrière has shown that closed $O(n-1, 1)$ affine manifolds are complete. It is obvious that a $Sim(n, 0)$ affine manifold is incomplete, since an element of its holonomy which doesn't lie in $O(n, 0)$ fixes an element of $\mathbb{R}^n$. There exist examples of complete $Sim(n-1, 1)$ affine manifolds. Let's give one:

Endow $\mathbb{R}^n$ with its basis $(e_1, \dots, e_n)$ and with the lorentzian product defined by

$$\langle e_i, e_i \rangle_L = 1; 0 < i < n; \langle e_i, e_j \rangle_L = 0; i \neq j; \langle e_n, e_n \rangle_L = -1.$$

We restrict this product to $\mathbb{R}^2$. The affine map whose linear part is

$$\begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}$$

in the basis $(e_1 + e_2, e_1 - e_2)$, and whose translation part is $e_1 - e_2$ generates a group which acts properly and freely on $\mathbb{R}^2$.

The goal of this paper is to study closed $Sim(n-1, 1)$ affine manifolds. First we show:

Theorem 1. *A compact $Sim(n-1, 1)$ affine manifold is incomplete.*

After, using the notion of discompacity, we show

Theorem 2. *Let M be a closed radiant $Sim(n-1, 1)$ affine manifold whose developing map is injective, if a connected component C of $\mathbb{R}^n - q^{-1}(0)$ intersects $D(M)$, then either C is contained in $D(M)$ or a connected component of $C - H$, where H is an hyperplane.*

Interesting structures of $Sim(n-1, 1)$ affine manifolds can be constructed using the work of Goldman on projective structures on surfaces see [Gol]. For instance a $Sim(2, 1)$ structure which linear holonomy is Zariski dense in $Gl(3, \mathbb{R})$ is given in this paper.

1. Closed $Sim(n-1, 1)$ affine manifolds are incomplete.

The main goal of this part is to show that a closed $Sim(n-1, 1)$ affine manifold cannot be complete.

Let suppose that there exists a complete closed $Sim(n-1, 1)$ affine manifold M ; M is the quotient of $\mathbb{R}^n$ by a subgroup of affine transformations Γ , whose linear part is contained in $Sim(n-1, 1)$.

Lemma 1.1. *Let γ be an element of Γ whose linear part has a determinant < 1 . Then there exists a basis $(e_1, \dots, e_n)$ of $\mathbb{R}^n$ such that the linear part of γ in this basis has the following form:*

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{\lambda^2} & 0 \\ 0 & 0 & \frac{1}{\lambda} B'' \end{pmatrix}$$

where λ is a real number strictly superior to 1 in absolute value, and B'' is a matrix which preserves the restriction of an euclidean product to the sub vector space generated by $e_3, \dots, e_n$.

Proof.

We have supposed that the determinant of the linear part $L(\gamma)$ of γ , is strictly inferior to 1 in absolute value. This implies that there exists a real number $\lambda > 1$ such that $\lambda L(\gamma) = L(\gamma)'$, where $L(\gamma)'$ is an element of $O(n-1, 1)$. The linear map $L(\gamma)$ has 1 as eigenvalue, since γ acts freely. We deduce that λ is an eigenvalue of $L(\gamma)'$. We remark that $L(\gamma)'$ has another eigenvalue α which module is different from 1 and the module of λ since the absolute value of its determinant is 1. If α is not a real number, then α and its complex conjugated $\bar{\alpha}$ are eigenvalues associated to the complex eigenvectors u_1 and u_2 . In this case the restriction of $L(\gamma)'$ to the plane generated by $u_1 + u_2$ and $i(u_1 - u_2)$ is an euclidean similitude whose ratio is different from 1. This is impossible since $L(\gamma)'$ lies in $O(n-1, 1)$. Let v_1 and v_2 be the eigenvectors associated to λ and α and $\langle, \rangle_L$ the lorentzian product preserved by the linear holonomy. We have:

$$\langle v_1, v_1 \rangle_L = \langle v_2, v_2 \rangle_L = 0.$$

We deduce that the restriction of $\langle, \rangle_L$ to the plane P generated by v_1 and v_2 is nondegenerate and has signature $(1, 1)$. This implies that the restriction of $\langle, \rangle_L$ to the orthogonal W of P with respect to itself is a scalar product. The restriction B'' of $L(\gamma)'$ to W is an orthogonal linear map. We can suppose that its determinant is 1. We deduce that $\alpha = \frac{1}{\lambda}$.

Up to a change of origin, we can suppose that $\gamma(0) = (a_1, 0, \dots, 0)$ where a_1 is a real number. The restriction B of $L(\gamma)$ to the linear subspace generated by $(e_2, \dots, e_n)$ is strictly contracting. It is easy to show that the group generated by γ is not cocompact, so Γ contains another element γ_1 different from γ .

Lemma 1.2. *Let C be the linear part of γ_1 , then $C(e_1) = e_1 + b$ where b lies in the linear subspace generated by $e_2, \dots, e_n$.*

Proof.

Let k be an element of $\mathbb{N}$. Consider the element $\gamma^k \gamma_1$. Its linear part has 1 as eigenvalue. The matrix of this linear part in the basis $(e_1, \dots, e_n)$ is $A^k C$, where A is the matrix of the linear part of γ . Let u_k be an eigenvector of $A^k C$ associated to 1. We assume that the norm of u_k with respect to the euclidean scalar product defined $\langle e_i, e_j \rangle = \delta_{ij}$ is 1. Let $u_k = (u_k^1, u_k^2)$. We have $C(u_k) = (v_k^1, v_k^2)$, where u_k^1 and v_k^1 are elements of $\mathbb{R}$, and u_k^2 and v_k^2 are elements of the vector space F generated by $e_2, \dots, e_n$. We have $v_k^1 = u_k^1$ since $A^k(e_1) = e_1$, and A^k preserves F . Since B is strictly contracting the norm of $A^k(0, v_k^2)$ goes to 0 with respect to the euclidean norm. So u_k goes to e_1 and $C(e_1)$, which is the limit of $C(u_k) = A^{-k}(u_k)$ is $e_1 + b$ where b is an element of F .

First proof of the theorem 1.

Let c be the translational part of γ_1 in the basis $(e_1, \dots, e_n)$. Put $c = (c_1, \dots, c_n)$. We have $\gamma^k \circ \gamma_1 \circ \gamma^{-k}(0) = (c_1, B^k(-ka_1b + (c_2, \dots, c_n)))$. Since B is contracting and the action of Γ is proper, we deduce that $(c_2, \dots, c_n) = b = 0$. This implies that $\gamma^m \gamma_1^n(0) = (na_1 + mc_1)e_1$. Since the action of Γ on $\mathbb{R}^n$ is

proper and free we deduce that the subgroup $\{n, m \in \mathbb{Z}, na_1 + mc_1\}$ is discrete. We deduce from this fact that there exist p, q in $\mathbb{Z}$ such that $pa_1 + qc_1 = 0$. This implies that $\gamma^p = \gamma_1^{-q}$.

Let K be a fundamental domain of the action of Γ , recall that it is a compact such that for each $x \in \mathbb{R}^n$, there exists an element γ_x of Γ such that $\gamma_x(x) \in K$, and for each element γ of Γ , $\gamma(K^0) \cap K^0$ is empty, where K^0 is the interior of K . Let $\| \cdot \|$ be a norm associated to $\langle \cdot, \cdot \rangle$, the scalar product for which $(e_1, \dots, e_n)$ is an orthonormal basis. There exists a real $A > 0$ and an integer l such that $\|K\| < A$ and $l \mid a_1 \mid > A$. Consider the element $u = (0, y_0)$ of $\mathbb{R}^n$ where y_0 is an element in the vector subspace generated by $(e_3, \dots, e_n)$ if $n \geq 3$, otherwise y_0 is an element of $\text{Vect}(e_2)$ such that $\|B^l(y_0)\| > A$. There must exist an element γ_0 in Γ such that $\gamma_0(u)$ is in K . We have shown that there exist elements p, q in $\mathbb{Z}$ such that $\gamma_0^p = \gamma^q$. Denote by D the restriction of γ_0 to $\text{Vect}(e_2, \dots, e_n)$, and d_1 the real number such that $\gamma_0(0) = d_1 e_1$. We have $D^p = B^q$ and $pd_1 = qa_1$. Since $\|y_0\| > A$, D is contractant and $q > p$. It results from lemma 1.1 that the restriction of D and B to $\text{Vect}(e_3, \dots, e_n)$ are similarities of respective ratio r_d and r_b (If the dimension is 2, we consider the restriction of D and B to $\text{Vect}(e_2)$ which is a similarity). We have $(r_d)^p = (r_b)^q$. Moreover $\|B^l(y_0)\| = (r_b)^l \|y_0\| > A$, and $A > \|D(y_0)\| = r_d \|y_0\|$. We deduce that $(r_b)^l > r_d$. This implies that $(r_b)^{lq} > (r_d)^q$, which is equivalent to saying that $(r_b)^{lq} = (r_d)^{lp} > (r_d)^q$. This implies that $q > lp$.

p and q are elements of $\mathbb{N}$. We have that $\gamma_0(u) = (d_1, D(y_0)) \in K$. This implies that $A > |d_1|$. But we have $qA > lpA > lp \mid d_1 \mid = lq \mid a_1 \mid$. Which is a contradiction since we have supposed that $l \mid a_1 \mid > A$.

We can also deduce the proof of Theorem 1, from this deep result

Theorem [F-G-H].

Let M a compact affine manifold, which is the quotient of $\mathbb{R}^n$ by the group Γ which acts properly and freely; then Γ does not preserve proper affine subspaces.

Proof. Up to a finite cover, we can assume that M is oriented. We denote by $(C(M), d)$ the simplicial complex we define the simplicial homology of M . We can lift the simplicial decomposition to $\mathbb{R}^n$, and thus lift the simplicial complex $(C(M), d)$ to the simplicial complex $(C(\mathbb{R}^n, d'))$. This last complex has a structure of a $\mathbb{Z}\Gamma$ module. Its homology is trivial since $\mathbb{R}^n$ is contractible. We deduce that it is a resolution of Γ . We can use this resolution to calculate the real cohomology of Γ . The cohomology obtained is also the real cohomology of M . The De Rham theorem implies that $H^n_{DR}(M, \mathbb{R}) = H^n(M, \mathbb{R}) = H^n(\Gamma, \mathbb{R}) = \mathbb{R}$ since we have supposed that M is oriented. Suppose that Γ preserves an l -affine subspace F of $\mathbb{R}^n$. We denote by N the quotient of F by Γ . We have also that $H^n_{DR}(N, \mathbb{R}) = H^n(\Gamma, \mathbb{R}) = \mathbb{R}$. Since N is compact, it implies that $l \geq n$, we deduce that $l = n$.

Second proof of theorem 1.

Let c be the translational part of γ_1 as in the first proof, we can remark that in the basis $(e_1, \dots, e_n)$, we have $c = (c_1, 0, \dots, 0)$ this implies that Γ preserves the line $\mathbb{R}e_1$. This fact contradicts the previous theorem.

Remark.

The both proofs of theorem 1 are related. While proving theorem 1, we have shown that for every element γ_1 in Γ , there exists p , and q in $\mathbb{Z}$ such that $\gamma_1^p = \gamma_q$ this implies that the quotient of Γ by the group generated by Γ is a torsion group. We deduce that the real cohomological dimension of Γ is one which is contrary to the fact that up to a finite cover $H^n(M, \mathbb{R}) = H^n(\Gamma, \mathbb{R}) = \mathbb{R}$.

In contrast to the $Sim(n, 0)$ affine manifolds (See [Fr] theorem 1), there exist compact $Sim(n - 1, 1)$ affine manifolds which are not radiant. Here is an example.

Endow $\mathbb{R}^2$ with the Lorentzian product $(,)$ such that $(e_1, e_1) = (e_2, e_2) = 0$, and $(e_1, e_2) = 1$.

Consider the subgroup Γ of $Aff(\mathbb{R}^2)$ generated by the following transformations:

$$\begin{aligned}\gamma_1(x, y) &= (x + 1, y), \\ \gamma_2(x, y) &= (x, 2y),\end{aligned}$$

the quotient of $\mathbb{R} \times (\mathbb{R} - \{0\})$ by Γ is a compact $Sim(n - 1, 1)$ affine manifold.

2. On the universal cover of compact $Sim(n - 1, 1)$ affine manifolds.

In this part we are going to find properties of the universal cover of a closed radiant $Sim(n - 1, 1)$ affine manifold. We use the notion of discompacity defined by Carrière [Car] 2.2.1. Let us recall it.

We consider in $\mathbb{R}^n$ the unit ball B_n . The euclidean metric induces on closed subsets of $\mathbb{R}^n$ the Hausdorff distance. Let G be a subgroup of $Gl(n, \mathbb{R})$, and $(g_p)_{p \in \mathbb{N}}$ a sequence of elements of G . The limit of the family $(g_p(B_n) \cap B_n)_{p \in \mathbb{N}}$ converges in B_n . It is a degenerated ellipsoid (see [Car]). The codimension of this ellipsoid is the discompacity d , of the family $(g_p)_{p \in \mathbb{N}}$, the discompacity of the group with respect to the euclidean metric is the smallest d .

Obviously we cannot use the notion of discompacity in this form since the linear holonomy of our manifold may contain homotheties. Denote $q : \mathbb{R}^n \rightarrow \mathbb{R}$ $x \rightarrow \langle x, x \rangle_L$. We can define in $\mathbb{R}^n - q^{-1}(0)$ the metric

$$(u, v) \longrightarrow \langle u, v \rangle' = \frac{\langle u, v \rangle_{euc}}{q(x)}$$

where u and v are vectors of the tangent space at x and $\langle, \rangle_{euc}$ is the euclidean scalar product.

Theorem 2.1.

Let $\hat{x}$ be an element of $\hat{M}$, u and v , elements of $T_{\hat{x}}\hat{M}$, such that the geodesics $c_1 : [0, 1] \rightarrow \hat{M}$, $t \rightarrow \exp_{\hat{x}}(tu)$, and the one $c_2 : [0, 1] \rightarrow \hat{M}$, $t \rightarrow \exp_{\hat{x}}(tv)$ are defined. Suppose that the elements $\exp_{\hat{x}}(u)$ and $\exp_{\hat{x}}(v)$ can't be joined by a geodesic, but for every $t, t' < 1$, there is a geodesic between $\exp_{\hat{x}}(tu)$ and $\exp_{\hat{x}}(t'v)$. Let $c : [0, 1] \rightarrow \mathbb{R}^n$, $t \rightarrow \exp_{D(\exp(u))}(tw)$ be the geodesic between $D(\exp_{\hat{x}}(u))$ and $D(\exp_{\hat{x}}(v))$, and let $U_{\hat{x}}$ be the domain of definition of $\exp_{\hat{x}}$. Consider the element $t_0 \in [0, 1]$ such that for every $t < t_0$, $\exp_{D(\exp(u))}(tw)$

is an element of $D(\exp_{\hat{x}}(U_{\hat{x}}))$ but not is $\exp_{D(\exp_{\hat{x}}(u))}(t_0 w) = y$. Then y is an element of $q^{-1}(0)$.

Proof.

There is a geodesic $\hat{c}_3 : [0, 1[\rightarrow \hat{M}$, $t \rightarrow \exp_{\hat{x}}(tb)$ such that y is an element of the adherence of $D(\hat{c}_3([0, 1[))$ and such that $D(\hat{c}_3)$ is contained in the convex hull of $D(\hat{c}_1)$ and $D(\hat{c}_2)$, where $\hat{c}_1$, and $\hat{c}_2$ are geodesics of $\hat{M}$ respectively above c_1 and c_2 . Set $p(\hat{x}) = x$, the image c_3 of $p(\hat{c}_3)$ is a maximal incomplete geodesic of M . Since M is compact, there exists an element z of M such that the geodesic c_3 is recurrent in an affine chart U which contains z . We deduce as Carrière, the existence of a family of ellipsoids s_p of $\mathbb{R}^n$ whose centers are elements of $D(\hat{c}_3)$, such that for each p, p' , there is an element $\gamma_{p,p'}$ of the holonomy such that $\gamma_{p,p'}(s_p) = s_{p'}$ and the centers x_p of s_p goes to y .

Suppose that y is not an element of $q^{-1}(0)$.

Let z_p be an element of an ellipsoid s_p , and u_p, v_p two vectors in its tangent space. Put $\gamma_{p,p'} = \lambda_{p,p'} g_{p,p'}$ where $g_{p,p'}$ is an element of $O(n-1, 1)$. We have:

$$\frac{\langle \gamma_{p,p'}(u_p), \gamma_{p,p'}(v_p) \rangle_{euc}}{q(\gamma_{p,p'}(x))} = \frac{\langle g_{p,p'}(u_p), g_{p,p'}(v_p) \rangle_{euc}}{q(x)}.$$

Since the holonomy of M is supposed to be radiant.

The metrics $\langle, \rangle_{euc}$ and $\langle, \rangle'$ are equivalent in a neighborhood of y since $q(y)$ is different from 0. We know that the discompacity of the family of g_p in respect to the riemannian metric $\langle, \rangle_{euc}$ is 1. The family of ellipsoids s_p goes to an ellipsoid, or a codimension 1 degenerated ellipsoid centered in y . We conclude as in Carrière that y must be an element of $D(\exp_{\hat{x}}(U_{\hat{x}}))$. This is not possible, so $q(y) = 0$.

A similar result is given in [Gol].

Corollary 2.2. *Let M be a compact radiant $Sim(n-1, 1)$ affine manifold, let $\hat{x}$, u , and v be respectively elements of $\hat{M}$ and $T_{\hat{x}}\hat{M}$, such that $\exp_{\hat{x}}(u)$ and $\exp_{\hat{x}}(v)$ are defined. If the convex hull E of $(D(\hat{x}), D(\exp_{\hat{x}}(u)), D(\exp_{\hat{x}}(v)))$ is contained in a connected component of $\mathbb{R}^n - q^{-1}(0)$, then it is contained in $D(\exp_{\hat{x}}(U_{\hat{x}}))$.*

Proof.

Suppose that E is not contained in $D(\exp_{\hat{x}}(U_{\hat{x}}))$. Let y and z be two elements of $E \cap D(\exp_{\hat{x}}(U_{\hat{x}}))$ such that $y = D(\exp_{\hat{x}}(u_1))$, $z = D(\exp_{\hat{x}}(u_2))$, and for every $t_1, t_2 < 1$, $\exp_{\hat{x}}$ is defined on the convex hull of $0, tu_1, tu_2$, but the elements $\exp_{\hat{x}}(u_1)$ and $\exp_{\hat{x}}(u_2)$ cannot be joined by a geodesic. Consider the geodesic $c : [0, 1] \rightarrow \mathbb{R}^n$, $t \rightarrow \exp_y(tw)$ between y and z . There exists a real number $0 < t_0 < 1$, such that for $0 < t < t_0$, $\exp_y(tw)$ lies in $D(\exp_{\hat{x}}(U_{\hat{x}}))$, but not $\exp_y(t_0 w)$. We deduce from the theorem 2.1. that $\exp_y(t_0 w)$ must lie in $q^{-1}(0)$. This is contrary to the hypothesis.

More generally, we can determine, how the boundary of the image of the developing map of a compact radiant $Sim(n-1, 1)$ affine manifold is: more precisely, we have the following proposition which implies theorem 2:

Proposition 2.3. *Let M be a compact radiant $Sim(n-1, 1)$ affine manifold whose developing map is injective, the boundary of $D(\hat{M})$, is contained in the union of $q^{-1}(0)$ and an hyperplane.*

Proof.

As in [Car] p. 625, one can remark that elements of the boundary of $D(\hat{M})$ which are not elements of $q^{-1}(0)$ are limits of $(\gamma_n e)_{n \in \mathbb{N}}$, where γ_n is an element of the holonomy and e is an ellipsoid. We conclude that those elements are contained in at most two hyperplane H_1, H_2 . The case of two hyperplane is impossible, since those hyperplane are stable by the holonomy, the affine function α such that $\alpha(H_1) = 0$ and $\alpha(H_2) = 1$, will be invariant by the holonomy and so define a differentiable function on M without maximal. (It is the same argument used in [Car]).

Proposition 2.4. *Let M be a compact radiant affine manifold, if the image of the developing map is a convex set contained in an open set of $\mathbb{R}^n - q^{-1}(0)$, then the developing map is injective.*

Proof.

Let $\hat{x}$ be an element of $\hat{M}$. For every elements, u and v of $U_{\hat{x}}$, the convex hull of $D(\hat{x})$, $y = D(\exp_{\hat{x}}(u))$ and $z = D(\exp_{\hat{x}}(v))$ is a subset of $D(\hat{M}) \cap (\mathbb{R}^n - q^{-1}(0))$. We deduce from the corollary 2.2 that y and z are elements of $D(U_{\hat{x}})$. This implies that $U_{\hat{x}}$ is a convex set. We can conclude by using [Kos].

A particular case of the situation of corollary 2.4 is the following: endow a compact oriented surface S of genus > 2 , with an hyperbolic structure, and consider q the Lorentzian form defined on $\mathbb{R}^3$ by $q(x_1, x_2, x_3) = x_1^2 + x_2^2 - x_3^2$. The hyperbolic structure can be defined by a representation of the fundamental group of S , $\pi_1(S) \rightarrow O(2, 1)$ such that the quotient of $H = q^{-1}(-1)$ by $\pi_1(S)$ is S . The quotient of $W = \{x : q(x) < 0, x_3 > 0\}$ by the group generated by $\pi_1(S)$, and a homothetie of ratio $0 < \lambda < 1$, is a compact $Sim(n-1, 1)$ affine manifold whose universal cover is W .

More generally we have

Corollary 2.5. *Let M be a radiant compact affine manifold such that the image of its developing map is contained in $W = \{x : q(x) < 0, z > 0\}$, M is the quotient of a connected component of $W - H$ by a discrete group of $Sim(n-1, 1)$, where H is an hyperplane of $\mathbb{R}^n$.*

Proof. We remark that the interior of a connected component of $W - H$ is convex. This implies the image of the developing map is a convex set. The result follows using 2.3 and 2.4.

Let M be a compact radiant $Sim(n-1, 1)$ affine manifold, the foliation $D(\hat{\mathcal{F}}_q)$ of $\mathbb{R}^n - \{0\}$ whose leaves are the sub manifolds defined by $q = \text{constant}$ is invariant by the holonomy of M . Its pull back on $\hat{M}$ defines a foliation $\hat{\mathcal{F}}_q$ of $\hat{M}$, which gives rise to a foliation $\mathcal{F}_q$ of M . If $D(\hat{N}) = D(\hat{M}) \cap q^{-1}(0)$ is not empty, then $N = p(D^{-1}(D(\hat{N})))$ is a compact submanifold of M . Note that the 1-parameter group ϕ_t generated by the radiant vector field X_0 , preserves

the foliation $\mathcal{F}_q$, and is transverse to all the leaves but not to the connected components of N .

Proposition 2.6. *If N is empty, then M is the total space of a bundle over S^1 .*

Proof.

If N is empty, then ϕ_t is transverse to the foliation $\mathcal{F}_q$. This implies that this foliation is a Lie foliation. We conclude using [God] Corollary 2.6 p. 154.

Remark.

Recall that a riemannian foliation on a manifold M , is a foliation $\mathcal{F}$ of M such that there exists a riemannian metric of M which projects locally along the leaves of M , this is equivalent to saying that locally the distance between the leaves is well-defined, or the foliation is defined by locally submersions $M \rightarrow N$ which transitions functions preserve a riemannian metric of N .

Suppose that the image of the developing map is included in the upper cone $C = \{x/q(x) < 0, x_n > 0\}$, one can define a map $f : C \rightarrow q^{-1}(-1)$ such that $f(x)$ is the element of C colinear to x such that $q(f(x)) = -1$. Let $(\cdot, \cdot)_L$ be the flat Lorentzian product of $\mathbb{R}^n$, The restriction of the lorentzian product defined by $(u, v)_x = \frac{(u, v)_L x}{(x, x)_L}$ to $q^{-1}(-1)$ is an hyperbolic metric. This endows the radial flow $\hat{\phi}_t$ of C with a transverse riemannian structure. Since the holonomy of the manifold is included in $RO(n-1, 1)$, the riemannian structure of $\hat{\phi}_t$ pushes forward to a riemannian structure of its radial flow ϕ_t . In fact the flow ϕ_t is also a transversally $(O(n-1, 1), q^{-1}(-1))$ homogeneous foliation where $O(n-1, 1)$. This enables one to define the global holonomy h_ϕ of ϕ_t which is a representation $h_\phi : \pi_1(M) \rightarrow O(n-1, 1)$. It assigns to any element γ of $\pi_1(M)$ (identified in this case as a subgroup of $RO(n-1, 1)$), the element $t\gamma (t \in \mathbb{R})$ such that $t\gamma$ is an element of $O(n-1, 1)$ and it preserves C .

The riemannian foliation have been intensively studied, It has been shown by Molino that the adherence of a leaf of a riemannian foliation defined on a compact manifold is a submanifold. Carriere and Carron have shown that in the case of riemannian flows, the adherence of leaves are torus.

More precisely, for a riemannian foliation, let denote by $(M_1, \mathcal{F}_1)$ the bundle of transverse orthogonal frames of M endowed with the pulls back $\mathcal{F}_1$ of $\mathcal{F}$, one can define the sheaf of local vector fields of M_1 which commute with the global foliated vector fields of $\mathcal{F}_1$. This sheaf pushes forward to a sheaf $C(M, \mathcal{F})$ of M . It is called the commuting sheaf of $\mathcal{F}$. The second structure theorem of riemannian foliations says that the adherences of the leaves of $\mathcal{F}$ are orbits of the pseudogroup defined by $C(M, \mathcal{F})$. In the case of a riemannian flow, the Lie algebra $C(M, \mathcal{F})$ is commutative.

In the other hand consider the adherence L of the image of h_ϕ in $O(n-1, 1)$. If the image of h_ϕ is discrete, it implies that the orbit of ϕ_t are closed, the holonomy of a leaf of ϕ_t is finite, this implies that up to a finite cover we can consider that the foliation ϕ_t does not have holonomy. This finite cover of M is the total space of a bundle whose typical fiber is an hyperbolic manifold. This manifold is the quotient of $q^{-1}(-1)$ by the image of h_ϕ .

If L is not discrete, then its lie algebra l is isomorphic to $C(M, \mathcal{F})$ see [W], we deduce that this Lie algebra is commutative. This imply that the connected component L_0 of L is a non trivial commutative group. Since $\pi_1(M)$ normalizes L_0 , using [G-K] 1.3 one can conclude that $\pi_1(M)$ is a solvable group. We have shown:

Theorem.

Suppose that the image of the developing map of a $Sim(n-1, 1)$ compact radiant affine manifold is contained in $C = \{x \in \mathbb{R}^n, 0 > q(x), \text{ and } x_n > 0\}$, then if $\pi_1(M)$ is not solvable, the leaves of the radiant flow are compact.

Remark.

The previous result is a particular case of a result due to Epstein for transversely hyperbolic foliation. Moreover if Epstein shows that if the leaves of the radiant flow are not compact, then the dimension of M is 3 or 4, and M is a the quotient of a solvable group G endowed with a left symmetric structure.

References.

- [Car] Carrière, Y. Autour de la conjecture de L. Markus sur les variétés affines. Invent. math. 95, (1989) 615-628.
- [Car-Car] Caron, P. Carrière, Y. Flots transversalement de Lie $\mathbb{R}^n$, flots de Lie transversalement minimaux
- [E] Epstein, D. Transversally hyperbolic 1-dimensional foliation Asterisque 116.
- [Fr] Fried, D. Closed similarity manifolds. Comment. Math. Helvetici 55, (1980) 576-582.
- [F-G-H] Fried, D. Goldman, W. Hirsch, M. Affine manifolds with nilpotent holonomy. Comment. Math. Helvetici 56, (1981) 487-523.
- [God] Godbillon, C. Feuilletages, études géométriques, I. Progress. in Math. vol 98. (1991).
- [Gol] Goldman, W. Projective structures with Fuchsian holonomy. J. Differential Geometry 25, (1987) 297-326.
- [G-K] Goldman, W. Kamishima, Y. The fundamental group of a compact flat Lorentz space form is virtually polycyclic. J. Diff. Geom. 19 (1984) 233-240.
- [Kos] Koszul, J-L. Variétés localement plates et convexité. Osaka J. Math. 2, (1965) 285-290.
- [L] Leslie, J. A remark on the group of automorphisms of a foliation having dense leaf. Jour. of Diff. Geom., 7 (1972) 597-601
- [Mo] Molino, P. Riemannian foliation, Progress in Mathematics (1988)
- [W] Wolak, R. Foliated G -structures and Riemannian foliations, Manuscripta. Math. 66 (1989), 45-59.